

Name _____
(Please Print)Chem 104 - Section 1
Sample Final Examination

This test consists of eleven (11) pages, including this cover page. Be sure your copy is complete before beginning your work. If this test packet is defective, ask for another one.

The following tables, which may be used for any problem, will be distributed separately: periodic table; conjugate acid-base pairs and K_a 's; oxidizing and reducing agents and E° values.

Give all numerical answers to the proper number of significant figures.

You must show work leading to your numeric answers, except in question 1.

$$R = 0.0821 \text{ L}\cdot\text{atm}/\text{K}\cdot\text{mol} = 8.314 \text{ J}/\text{K}\cdot\text{mol}$$

$$1 \text{ Faraday} = 9.65 \times 10^4 \text{ Coulombs}$$

$$E_{\text{cell}} = E^\circ_{\text{cell}} - (0.0592/n)\log Q \text{ at } 25^\circ\text{C}$$

$$\text{K} = ^\circ\text{C} + 273$$

$$K_w = 1.00 \times 10^{-14} \text{ at } 25^\circ\text{C}$$

$$1.00 \text{ mol ideal gas} = 22.4 \text{ L at STP}$$

DO NOT WRITE BELOW THIS LINE

This is a copy of a typical final exam given in Chem 104. Your test will be different. This test is being posted to give you a sense of the format, style, scope, and level of a typical final exam in this course. This test may have questions on topics that will not be covered on the test you take. Moreover, your test may have questions on topics that are not covered on this test. Posting this test in no way limits the format, style, scope, or level of the test that you will take. **Do not limit your preparation to the material on this sample test.**

Name _____

1. (100 points; 4 points each) Circle the correct answers.

a. Has dipole-dipole intermolecular forces

NaCl C₆H₆ NH₃ I₂ SiO₂

b. Would make an acidic solution in water

NaHCO₃ (CH₃)₃N KNO₃ Fe(NO₃)₃ NaC₆H₅CO₂

c. Does not have hydrogen bonding

H₂O NH₃ CH₄ C₆H₅OH HF

d. "At constant pressure, the volume of a gas sample is proportional to its absolute temperature" is a statement of

Boyle's Law Gay-Lussac's Law Amonton's Law Dalton's Law Charles' Law

e. A 0.100 *m* solution of sugar in water has a freezing point of -0.186°C. Ideally, a 0.100 *m* solution of Na₃PO₄ should have a freezing point of

-0.186 °C -0.372 °C -0.558 °C -0.744 °C -0.062 °C

f. A 4.00-L sample of an ideal gas in a piston chamber at 200 K and 1.00 atm is heated to 300 K and the pressure is raised to 1.50 atm. The new volume will be

4.00 L 6.00 L 9.00 L 0.750 L 1.50 L

g. Consider the equilibrium C₂H₄(g) + H₂(g) ⇌ C₂H₆(g) in a container of fixed volume. Increasing the total pressure by introducing some inert He(g) will cause

K_p to increase *K_p* to decrease more C₂H₆ (g) to form

more C₂H₄(g) + H₂(g) to form no change

[Continue to the next page.]

Name _____

- h. The equilibrium $A_2(g) \rightleftharpoons 2A(g)$ is a single-step reaction in both directions. For the forward reaction $A_2(g) \rightarrow 2A(g)$ the activation energy E_a^f is 289 kJ and ΔH is +125 kJ. Therefore, E_a^r , the activation energy for the reverse reaction $2A(g) \rightarrow A_2(g)$ is

– 125 kJ –289 kJ +164 kJ +414 kJ +125 kJ

- i. For the equilibrium $A_2(g) \rightleftharpoons 2A(g)$, described in part h, if the temperature were increased

[A] would increase, $[A_2]$ would decrease, and K_c would remain unchanged.

$[A_2]$ would increase, [A] would decrease, and K_c would remain unchanged.

[A] would increase, $[A_2]$ would decrease, and K_c would increase.

$[A_2]$ would increase, [A] would decrease, and K_c would decrease.

$[A_2]$, [A], and K_c would remain unchanged.

- j. Consider the equilibrium $A_2(g) \rightleftharpoons 2A(g)$, described in part h, for which $K_c = 44.2$ mol/L at 200 °C. If an equilibrium mixture at 200 °C is found to contain 0.100 mol/L of A(g), what is the concentration of $A_2(g)$ in mol/L?

0.442 5.00×10^{-2} 1.00×10^{-2} 9.05×10^{-4} 2.26×10^{-4}

- k. As the mechanism in part h indicates, the forward reaction $A_2(g) \rightarrow 2A(g)$ should obey the differential rate law $rate_f = k_f[A_2]$. Therefore, a plot of $\ln [A_2]$ versus time should give a straight line whose slope is

$-E_a/R$ $\ln [A]$ $t_{1/2}$ $-k_f$ $\ln [A_2]_0$

- l. The half-life of the forward reaction $A_2(g) \rightarrow 2A(g)$ described in part h is $t_{1/2} = 2.5$ s. If 1.6 mol $A_2(g)$ is introduced into an empty one-liter vessel at 200 °C, what will be the concentration of $A_2(g)$ in mol/L after the reaction has run for 10.0 s? [Assume the reverse reaction $2A(g) \rightarrow A_2(g)$ has been suppressed by removing product as it is formed.]

1.6 0.80 0.40 0.20 0.10

[Continue to the next page.]

Name _____

- m. What is the *molality* (m) of a solution prepared by dissolving 36.04 g of glucose (m.w. = 180.2 u) in 275 g of water?

0.727 0.655 0.200 7.25×10^{-4} 312

- n. What is the expected osmotic pressure in atmospheres (atm) at 298 K of a 0.250 M solution of $\text{CaCl}_2(aq)$?

6.12 12.2 18.3 6.05×10^2 1.82×10^3

- o. Consider a 0.10 M solution of the weak base B, for which $K_b = 1.0 \times 10^{-2}$. Which of the following expressions could be used to obtain a reasonably accurate value of the hydroxide concentration, $[\text{OH}^-]$, for this solution without carrying out unnecessary calculations?

$$[\text{OH}^-] = \sqrt{C_B K_b} \quad [\text{OH}^-] = C_B \quad K_b = \frac{[\text{OH}^-]^2}{C_B - [\text{OH}^-]}$$

$$K_b = K_w / K_a \quad \text{pOH} = \text{p}K_b + \log\left(\frac{C_{\text{BH}^+}}{C_B}\right)$$

- p. Which among the following is the strongest base in water?

F_3CCO_2^- $\text{C}_2\text{H}_5\text{OH}$ ClO_4^- CH_3CO_2^- HSO_4^-

- q. Nitrogen gas comprises 78% of air by volume and has a Henry's Law constant of 6.8×10^{-4} mol/L·atm for dissolving in water at 25 °C. What is the solubility at 25 °C in mol/L of nitrogen when the air pressure is 690 torr?

7.5×10^{-4} 6.8×10^{-4} 6.2×10^{-4} 5.3×10^{-4} 4.8×10^{-4}

- r. In KHSO_3 , the oxidation number for the S atom is


+6 +5 +4 -1 -2

[Continue to the next page.]

Name _____

[Consult the Table of Oxidizing and Reducing Agents for the following three questions.]

- s. Consider the reaction,



Under standard conditions this reaction is

spontaneous nonspontaneous at equilibrium




- t. Among the following, the strongest
- oxidizing agent*
- is



- u. Among the following, which one could be used to reduce
- Pb^{2+}
- to
- Pb^0
- (
- $E^\circ = -0.126 \text{ v}$
-) under standard conditions?



- v. The reaction
- $\text{X}_2(g) + \text{Y}_2(g) \rightarrow 2\text{XY}(g)$
- has the experimental rate law
- $\text{Rate} = k[\text{X}_2]$
- . Which one of the following four proposed mechanisms is consistent with this observed rate law?



[Continue to the next page.]

Name _____

The next three questions pertain to following rate data for the reaction $A_2 + B_2 \rightarrow 2AB$. (All concentrations are in mol/liter.)

	$[A_2]$	$[B_2]$	Initial rate (mol/L)·s ⁻¹
Experiment 1	0.15	0.15	2.0×10^{-4}
Experiment 2	0.15	0.60	8.0×10^{-4}
Experiment 3	0.60	0.15	4.0×10^{-4}

- w. In the differential rate law for this reaction, the dependence of rate on $[A_2]$ would be expressed as $Rate \propto$

$$[A_2]^0 \quad [A_2] \quad [A_2]^2 \quad [A_2]^{1/2} \quad [A_2]^{3/2}$$

- x. In the differential rate law for this reaction, the dependence of rate on $[B_2]$ would be expressed as $Rate \propto$

$$[B_2]^0 \quad [B_2] \quad [B_2]^2 \quad [B_2]^{1/2} \quad [B_2]^{3/2}$$

- y. The units on the rate constant, k , are

$$s^{-1} \quad (\text{mol/L}) \cdot s^{-1} \quad (\text{mol/L})^{-1} \cdot s^{-1} \quad (\text{mol/L})^{-2} \cdot s^{-1} \quad (\text{mol/L})^{-1/2} \cdot s^{-1}$$


[End of multiple choice questions.]

Name _____

2. (28 points) Given the following two couples and their corresponding standard reduction potentials:



Part I (20 points; 2 points each) Fill in the blanks below regarding the following galvanic cell under standard conditions.



⚠ *Do not assume that the cell is shown in the proper order (i.e., anode and cathode placed on the correct sides). You must determine whether or not it is properly shown.*

a. The oxidation half-reaction is

b. The reduction half-reaction is

c. The overall net ionic reaction for the cell is

d. $E^{\circ}_{\text{cell}} =$ _____ volts

e. The oxidizing agent is _____

f. The reducing agent is _____

g. In the cell diagram *as shown above*, the anode is shown on the _____ (right/left).

h. Is the cell diagram *as shown above* in conventional order? _____ (yes/no)

i. In the cell diagram *as shown above*, the direction of electron flow in the circuit wire is

_____ (left to right/right to left).


j. As the cell runs, E_{cell} will _____ (increase/decrease/stay the same).

[Part II on the following page]

Name _____

2. Part II (8 points) Calculate E_{cell} for the galvanic cell described in Part I (previous page) when $[\text{Cu}^{2+}] = 0.360 \text{ M}$ and $[\text{Pb}^{2+}] = 1.00 \times 10^{-3} \text{ M}$.

3. (16 points) Balance the following skeletal redox equation in *acidic solution* using the ion-electron (half-reaction) method:



In answering this question, you must show the following:


- (1) A balanced half-reaction for $\text{HClO}_2(aq) \rightarrow \text{Cl}^-(aq)$ in *acidic solution* (6 points);
- (2) A balanced half-reaction for $\text{As}(s) \rightarrow \text{HAsO}_2(aq)$ in *acidic solution* (6 points);
- (3) The balanced overall redox reaction, with lowest whole-number coefficients (4 points).

Don't bother writing the state designations (*aq*) and (*s*), **but be sure that charges on all species are properly shown.**

Name _____

4. (10 points) How many grams of nickel metal (at. wt. 58.71 u) will be produced at the electrode in the electrolysis of a $\text{Ni}(\text{NO}_3)_2$ solution for 8.00 amp for a period of 96.0 minutes? The half-reaction at the electrode is $\text{Ni}^{2+}(\text{aq}) + 2e^- \rightarrow \text{Ni}(\text{s})$.


5. (10 points) Consider the following equilibrium for which $K_c = 7.00$ at 400 K.



Calculate the equilibrium concentrations of all three species that should result when 2.50 mol of $\text{BrCl}(\text{g})$ are introduced into an empty one-liter vessel at 400 K.

Name _____

6. (18 points) Formic acid, HCO_2H , has $K_a = 1.9 \times 10^{-4}$ for its hydrolysis



- a. (6 points) What is the pH of 0.10 M solution of formic acid?
- b. (3 points) What is the pH of a solution in which the concentration of formic acid is 0.10 M and concentration of sodium formate is 0.10 M?
- c. (9 points) What is the pH of a 0.10 M sodium formate solution?

Name _____

7. (18 points) Consider the reaction



Given the following thermodynamic data:

Substance	ΔH_f° (kJ/mol)	ΔG_f° (kJ/mol)	S° (J/K·mol)
$\text{PCl}_5(\text{g})$	-398.9	-324.6	352.7
$\text{PCl}_3(\text{g})$	-306.4	-286.3	311.7
$\text{Cl}_2(\text{g})$	0	0	223.0

- a. (3 points) What is the value of ΔG° for the reaction at 25 °C? Is the reaction spontaneous at this temperature?
- b. (6 points) What is the value of K_p for the reaction at 25 °C?
- c. (9 points) Assuming ΔH and ΔS are constant with changes in temperature, calculate the value of ΔG for the reaction at 300 °C. Is the reaction spontaneous at this temperature?