

Chem 116 - Test 1 Practice Problems

1. Circle the best answer to each of the following.

a. If c is a constant, the equation that is the basis for Avogadro's hypothesis is

$$V = c/P \quad V = cn \quad P = cT \quad KE = cT \quad V = cT$$

b. A gas sample initially at 1.00 atm is expanded at constant temperature from 50.0 L to 75.0 L. the final pressure is

$$3.33 \text{ atm} \quad 1.50 \text{ atm} \quad 1.00 \text{ atm} \quad 0.667 \text{ atm} \quad 0.200 \text{ atm}$$

c. At STP a 14.3-g sample of gas occupies 5.00 L. What is its molecular weight?

$$2.86 \quad 14.3 \quad 22.4 \quad 64.1 \quad 112$$

d. In a gas mixture of He, Ne, and Ar with a total pressure of 8.40 atm, the partial pressures of He and Ne are 1.50 atm and 2.00 atm, respectively. What is the mole fraction of Ar in the mixture?

$$0.179 \quad 0.714 \quad 0.238 \quad 0.417 \quad 0.583$$

e. A 0.100-mole sample of oxygen gas (m.w. = 32.0) effused through a pin hole in 5.00 seconds. Under the same conditions, how long would it take the same amount of CO_2 (m.w. = 44.0) to effuse?

$$1.17 \text{ s} \quad 3.64 \text{ s} \quad 4.26 \text{ s} \quad 5.86 \text{ s} \quad 6.88 \text{ s}$$

f. Of the following gases, which would deviate most from ideal behavior?




g. Which of the following is *least* soluble in methanol, CH_3OH ?



h. Which of the following solutions would have the highest osmotic pressure?



i. Which of the following has the highest boiling point?



2. (26 points) Propylene glycol [$C_3H_6(OH)_2$, m.w. = 76.09 u] is sometimes used as a non-toxic alternative to the usual ethylene glycol automotive antifreeze. In addition to protecting against freezing in cold weather, a good antifreeze must also prevent boil-over of the engine coolant in hot weather. Consider a certain antifreeze solution that is 29.0 % by weight propylene glycol in water.

a. What is the molality of the solution?

b. At what temperature in $^{\circ}C$ will the solution boil? $K_b = 0.512 \text{ }^{\circ}C/m$ for water.

c. What is the mole fraction of *water* (not propylene glycol) in this solution? [m.w. $H_2O = 18.02$ u]

d. Assuming that the vapor pressure of the solution is entirely from water (i.e., propylene glycol is virtually nonvolatile), what is the vapor pressure above the solution in torr at water's normal boiling point of $100.0 \text{ }^{\circ}C$?

3. A 3.567-L sample of $\text{CO}_2(g)$ (m.w. = 44.01 u) is collected over water 35.40°C . The pressure inside the vessel is 772.2 torr. At 35.40°C the vapor pressure of water is 43.12 torr.

a. How many moles of $\text{CO}_2(g)$ does the sample contain?

b. What are the mole fractions of $\text{CO}_2(g)$ and $\text{H}_2\text{O}(g)$ in the sample?

4. A solution prepared by dissolving 0.525 g of an unknown non-electrolyte in enough water to make 125 mL of solution has an osmotic pressure of 1.10 atm at 27°C . What is the molar mass of the solute?